

Generic positivity of the Beilinson–Bloch height (joint with Shouwu Zhang)

Ziyang Gao

University of California, Los Angeles

Fort Collins, Colorado, SRI 2025

Motivation: Weil height

A. Weil (1928) defined **height** to measure the “size” of algebraic points.

- ✎ On \mathbb{Q} : $h(a/b) = \log \max\{|a|, |b|\}$, for $a, b \in \mathbb{Z}$ and $\gcd(a, b) = 1$.
- ✎ On $\mathbb{P}^n(\mathbb{Q})$: $h([x_0 : \cdots : x_n]) = \log \max\{|x_0|, \dots, |x_n|\}$, for $x_i \in \mathbb{Z}$ and $\gcd(x_0, \dots, x_n) = 1$.
- ✎ Arbitrary number field K : For $[x_0 : \cdots : x_n] \in \mathbb{P}^n(K)$,

$$h([x_0 : \cdots : x_n]) = \frac{1}{[K : \mathbb{Q}]} \sum_{v \in M_K} \log \max\{\|x_0\|_v, \dots, \|x_n\|_v\}.$$

↪ (logarithmic) Weil height on $\mathbb{P}^n(\overline{\mathbb{Q}})$.



Motivation: Weil height

Two important properties →



Positivity

$h(\mathbf{x}) \geq 0$ for all $\mathbf{x} \in \mathbb{P}^n(\overline{\mathbb{Q}})$.

Northcott Property (1949)

For all B and $d \geq 1$,

$\{\mathbf{x} \in \mathbb{P}^n(\overline{\mathbb{Q}}) : h(\mathbf{x}) \leq B, [\mathbb{Q}(\mathbf{x}) : \mathbb{Q}] \leq d\}$

is finite.

Motivation: (naive) Height Machine

X projective variety defined over a number field K_0 .

- X can be embedded into \mathbb{P}^N \rightsquigarrow naive height h_{Weil} on $X(\overline{\mathbb{Q}})$
- Different embeddings \rightsquigarrow well-defined up to a bounded function.

Two important properties \rightarrow



Bounded from below

There exists C such that $h_{\text{Weil}}(x) \geq C$ for all $x \in X(\overline{\mathbb{Q}})$.

Northcott Property

For all B and $d \geq 1$,

*$\{x \in X(\overline{\mathbb{Q}}) : h(x) \leq B, [K_0(x) : K_0] \leq d\}$
is finite.*

Motivation: Dominant height function

- X quasi-projective variety defined over $\overline{\mathbb{Q}}$;
- $h: X(\overline{\mathbb{Q}}) \rightarrow \mathbb{R}$.

Definition

h is called a *dominant height* if it has a lower bound and satisfies the Northcott property.

Two famous examples:

Example

Néron–Tate height on abelian variety A , with lower bound 0. \rightsquigarrow Mordell–Weil theorem, formulation of Birch and Swinnerton-Dyer Conjecture, etc.

Example (On the moduli space \mathbb{M}_g of smooth projective curves of genus g)

$h_{\text{Fal}}: \mathbb{M}_g(\overline{\mathbb{Q}}) \rightarrow \mathbb{R}$, sending each curve C to the Faltings height of its Jacobian.
 \rightsquigarrow Mordell Conjecture.

Motivation: Dominant height function

- X quasi-projective variety defined over $\overline{\mathbb{Q}}$;
- $h: X(\overline{\mathbb{Q}}) \rightarrow \mathbb{R}$.

Definition

h is called a *dominant height* if it has a lower bound and satisfies the Northcott property.

Two famous examples:

Example

Néron–Tate height on abelian variety A , with lower bound 0. \rightsquigarrow Mordell–Weil theorem, formulation of Birch and Swinnerton-Dyer Conjecture, etc.

Example (On the moduli space \mathbb{M}_g of smooth projective curves of genus g)

$h_{\text{Fal}}: \mathbb{M}_g(\overline{\mathbb{Q}}) \rightarrow \mathbb{R}$, sending each curve C to the Faltings height of its Jacobian.
 \rightsquigarrow Mordell Conjecture.

Motivation: Beilinson–Bloch height and conjecture

Aim (from 1980s):

- Extend height from points to higher cycles which are homologically trivial (Beilinson–Bloch height).
- Positivity of BB height.
- Finiteness of the rank of Chow group.
- Generalization of BSD.

Known results

- Conjecturally defined.
Unconditional in some cases (Gross–Schoen, Künnemann, S. Zhang).
- Some sporadic families.
- ???
- ???


Motivation: Gross–Schoen and Ceresa cycles

Example (BB height is known to be unconditionally defined)

- C smooth projective curve of genus $g \geq 2$;
- $\xi \in \text{Pic}^1(C)$ such that $(2g-2)\xi = \omega_C$.


From these data, we obtain homologically trivial 1-cycles:

- ✎ (Gross–Schoen) $\Delta_{\text{GS}}(C) \in \text{Ch}_1(C^3)$ the modified diagonal;
- ✎ (Ceresa) $\text{Ce}(C) := i_\xi(C) - [-1]^* i_\xi(C) \in \text{Ch}_1(J)$, with $J = \text{Jac}(C)$.



modified diagonal

$$\Delta_{123} - \Delta_{12} - \Delta_{23} - \Delta_{13} + \Delta_1 + \Delta_2 + \Delta_3.$$



Goal of the project

Propose a systematic way to study the positivity of the Beilinson–Bloch height $\langle \bullet, \bullet \rangle_{\text{BB}}$.

- Starting point: Use $\langle \bullet, \bullet \rangle_{\text{BB}}$ to define a function on a suitable parametrizing space.

Setup for our main result

Two functions on \mathbb{M}_g :

$$h_{\text{GS}}: \mathbb{M}_g(\overline{\mathbb{Q}}) \rightarrow \mathbb{R}, \quad [C] \mapsto \langle \Delta_{\text{GS}}(C), \Delta_{\text{GS}}(C) \rangle_{\text{BB}}$$

$$h_{\text{Ce}}: \mathbb{M}_g(\overline{\mathbb{Q}}) \rightarrow \mathbb{R}, \quad [C] \mapsto \langle \text{Ce}(C), \text{Ce}(C) \rangle_{\text{BB}}$$

- Facts:
- Both vanish on the hyperelliptic locus;
 - $h_{\text{GS}} = 6h_{\text{Ce}}$

Question (in different grades)

Assume $g \geq 3$.

- ✎ (i) Is h_{GS} a dominant height (*lower bound + Northcott property*) on a Zariski open dense subset U of \mathbb{M}_g defined over \mathbb{Q} ? \rightsquigarrow *generic positivity*
- ✎ (ii) Can we determine U ?
- ✎ (iii) Is the lower bound ≥ 0 ?

Our main result

Theorem (G'–S.Zhang, 2024)

Assume $g \geq 3$. Let $\mathbb{M}_g^{\text{amp}}$ be the maximal $\overline{\mathbb{Q}}$ -Zariski open subset of \mathbb{M}_g on which h_{GS} is a dominant height.

Then $\mathbb{M}_g^{\text{amp}}$ is non-empty and is defined over \mathbb{Q} . ✓ for (i)

Moreover, $\mathbb{M}_g^{\text{amp}}$ can be “constructed”. ✓ partially for (ii)

Still, we need to express $\mathbb{M}_g^{\text{amp}}$ more explicitly and need to show that the lower bound is ≥ 0 . But already, we have

Corollary (Generic positivity)

For any number field K , there are at most finitely many C/K lying in $\mathbb{M}_g^{\text{amp}}(\overline{\mathbb{Q}})$ such that $h_{\text{GS}}([C]) \leq 0$.

Key steps of our proof

Steps:

- h_{GS} defined by an a.l.b. $\overline{\mathcal{L}}$
- volume identity for $\text{vol}(\widetilde{\mathcal{L}})$

Bridged via:

- Algebraicity of Betti strata
- Non-vanishing of Betti form

Tools:

- Adelic line bundle (Yuan–Zhang 2021).
- Morse Inequality (Demailly 1991).

- ✎ Abel–Jacobi periods (Griffiths 1960s)
- ✎ archimedean local heights (Hain 1990s)

- Mixed Ax–Schanuel (Chiu/Gao–Klingler 2021).
- O-minimality for period map (Bakker, Brunebarbe, Klingler, Tsimerman 2018–2020...).

Adelic line bundle

Theorem

There exists an *adelic line bundle* $\overline{\mathcal{L}}$ on \mathbb{M}_g such that $h_{\text{GS}} = h_{\overline{\mathcal{L}}}$.

A construction was given by Yuan. We give a new construction using:

- Polarized dynamical system on the universal Jacobian $\text{Jac}(\mathcal{C}_g/\mathbb{M}_g) \rightarrow \mathbb{M}_g$.
- Deligne pairing to “push-forward” adelic line bundles on $\mathcal{C}_g \times_{\mathbb{M}_g} \mathcal{C}_g$ to \mathbb{M}_g .
- Explicit computation.
- We use our construction to prove the volume identity.

Adelic line bundle

✍ What is an adelic line bundle, and what is the motivation/idea behind?

Let (X, L) projective variety with a line bundle, defined over a number field K .

- Naive height $h_L: X(\overline{\mathbb{Q}}) \rightarrow \mathbb{R}$, *well-defined up to a bounded function*.
- **Wish to get genuine functions**. Sometimes okay, e.g. Néron–Tate height on abelian varieties.
- In general, **integral model** $(\mathcal{X}, \overline{\mathcal{L}})$, with $\overline{\mathcal{L}}$ a Hermitian line bundle.
But cannot recover Néron–Tate height in this way!!
- **Solution**: Put a \overline{K}_v -metric of L on $X(\overline{K}_v)$ for all $v \in M_K \rightsquigarrow$ **metrized line bundle**
An **adelic line bundle** $\overline{\mathcal{L}}$ is a metrized line bundle which can be obtained as a “limit” of integral models.
- This construction can be generalized to quasi-projective varieties, “limit” of integral models of compactifications of $X \rightsquigarrow$ generic fiber $\widetilde{\mathcal{L}}$ of $\overline{\mathcal{L}}$.

Example $(X = \text{Spec} K)$

An adelic line bundle on $\text{Spec} K$ is $(L, \{\|\cdot\|_v\}_v)$ with $L =$ vector space of dim 1 and $\|\cdot\|_v$ a K_v -metric, satisfying: $\forall \ell \in L \setminus \{0\}, \|\ell\|_v = 1$ for all but finitely many v .

Adelic line bundle

✍ What is an adelic line bundle, and what is the motivation/idea behind?

Let (X, L) projective variety with a line bundle, defined over a number field K .

- Naive height $h_L: X(\overline{\mathbb{Q}}) \rightarrow \mathbb{R}$, *well-defined up to a bounded function*.
- **Wish to get genuine functions**. Sometimes okay, e.g. Néron–Tate height on abelian varieties.
- In general, **integral model** $(\mathcal{X}, \overline{\mathcal{L}})$, with $\overline{\mathcal{L}}$ a Hermitian line bundle.
But cannot recover Néron–Tate height in this way!!
- **Solution**: Put a \overline{K}_v -metric of L on $X(\overline{K}_v)$ for all $v \in M_K \rightsquigarrow$ **metrized line bundle**
An **adelic line bundle** $\overline{\mathcal{L}}$ is a metrized line bundle which can be obtained as a “limit” of integral models.
- This construction can be generalized to quasi-projective varieties, “limit” of **integral models of compactifications of X** \rightsquigarrow generic fiber $\widetilde{\mathcal{L}}$ of $\overline{\mathcal{L}}$.

Example ($X = \text{Spec} K$)

An adelic line bundle on $\text{Spec} K$ is $(L, \{\|\cdot\|_v\}_v)$ with $L =$ vector space of dim 1 and $\|\cdot\|_v$ a K_v -metric, satisfying: $\forall \ell \in L \setminus \{0\}, \|\ell\|_v = 1$ for all but finitely many v .

Adelic line bundle

✍ What is an adelic line bundle, and what is the motivation/idea behind?

Let (X, L) projective variety with a line bundle, defined over a number field K .

- Naive height $h_L: X(\overline{\mathbb{Q}}) \rightarrow \mathbb{R}$, *well-defined up to a bounded function*.
- **Wish to get genuine functions**. Sometimes okay, e.g. Néron–Tate height on abelian varieties.
- In general, **integral model** $(\mathcal{X}, \overline{\mathcal{L}})$, with $\overline{\mathcal{L}}$ a Hermitian line bundle.
But cannot recover Néron–Tate height in this way!!
- **Solution**: Put a \overline{K}_v -metric of L on $X(\overline{K}_v)$ for all $v \in M_K \rightsquigarrow$ **metrized line bundle**
An **adelic line bundle** $\overline{\mathcal{L}}$ is a metrized line bundle which can be obtained as a “limit” of integral models.
- This construction can be generalized to quasi-projective varieties, “limit” of **integral models of compactifications of X** \rightsquigarrow generic fiber $\widetilde{\mathcal{L}}$ of $\overline{\mathcal{L}}$.

Example ($X = \text{Spec} K$)

An adelic line bundle on $\text{Spec} K$ is $(L, \{\|\cdot\|_v\}_v)$ with $L =$ vector space of dim 1 and $\|\cdot\|_v$ a K_v -metric, satisfying: $\forall \ell \in L \setminus \{0\}, \|\ell\|_v = 1$ for all but finitely many v .

Volume identity

Recall our main theorem

Theorem (G'–S.Zhang, 2024)

Assume $g \geq 3$. Then h_{GS} is a dominant height on a Zariski open dense subset $\mathbb{M}_g^{\text{amp}}$ of \mathbb{M}_g defined over \mathbb{Q} . ✓ for (i)

Moreover, $\mathbb{M}_g^{\text{amp}}$ can be “constructed”. ✓ partially for (ii)

➤ Part (i) except “defined over \mathbb{Q} ” $\Leftrightarrow \widetilde{\mathcal{L}}$ is big, i.e. $\text{vol}(\widetilde{\mathcal{L}}) > 0$.

A key property we prove is the following **volume identity**.

Theorem (GZ 2024)

$$\text{vol}(\widetilde{\mathcal{L}}) = \int_{\mathbb{M}_g(\mathbb{C})} c_1(\overline{\mathcal{L}})^{\wedge \dim \mathbb{M}_g}.$$

Stronger:
needed for
“over \mathbb{Q} ”

Theorem (GZ 2024)

For each subvariety S of $\mathbb{M}_{g,\mathbb{C}}$, we have

$$\text{vol}(\widetilde{\mathcal{L}}_{\mathbb{C}}|_S) = \int_{S(\mathbb{C})} c_1(\overline{\mathcal{L}})^{\wedge \dim S}.$$

Volume identity

Recall our main theorem

Theorem (G'–S.Zhang, 2024)

Assume $g \geq 3$. Then h_{GS} is a dominant height on a Zariski open dense subset $\mathbb{M}_g^{\text{amp}}$ of \mathbb{M}_g defined over \mathbb{Q} . ✓ for (i)

Moreover, $\mathbb{M}_g^{\text{amp}}$ can be “constructed”. ✓ partially for (ii)

➤ Part (i) except “defined over \mathbb{Q} ” $\Leftrightarrow \widetilde{\mathcal{L}}$ is big, i.e. $\text{vol}(\widetilde{\mathcal{L}}) > 0$.

A key property we prove is the following **volume identity**.

Theorem (GZ 2024)

$$\text{vol}(\widetilde{\mathcal{L}}) = \int_{\mathbb{M}_g(\mathbb{C})} c_1(\overline{\mathcal{L}})^{\wedge \dim \mathbb{M}_g}.$$

Stronger:
needed for
“over \mathbb{Q} ”

Theorem (GZ 2024)

For each subvariety S of $\mathbb{M}_{g,\mathbb{C}}$, we have

$$\text{vol}(\widetilde{\mathcal{L}}_{\mathbb{C}}|_S) = \int_{S(\mathbb{C})} c_1(\overline{\mathcal{L}})^{\wedge \dim S}.$$

Volume identity

Theorem (GZ 2024)

For each subvariety S of $\mathbb{M}_{g,\mathbb{C}}$, we have

$$\mathrm{vol}(\widetilde{\mathcal{L}}_{\mathbb{C}}|_S) = \int_{S(\mathbb{C})} c_1(\overline{\mathcal{L}})^{\wedge \dim S}.$$

- LHS defined using some kind of h^0 , so invariant under $\mathrm{Aut}(\mathbb{C})$.
↪ Used for “over \mathbb{Q} ” in the main theorem.
- In the flavor of (arithmetic) Hilbert–Samuel.
- **Problem:** $\widetilde{\mathcal{L}}$ is not known to be nef!!!
- **Solution:** Compute $\mathrm{vol}(\widetilde{\mathcal{L}}_{\mathbb{C}}|_S)$ directly, by our explicit construction of $\overline{\mathcal{L}} = \{(\mathcal{M}_i, \overline{\mathcal{L}}_i)\}_{i \geq 1}$ and the fact $\mathrm{vol}(\mathcal{L}_{i,\mathbb{Q}}|\overline{S}) \rightarrow \mathrm{vol}(\widetilde{\mathcal{L}}_{\mathbb{C}}|_S)$. Use **Demailly’s Morse Inequality** to bound $h^0(m\mathcal{L}_{i,\mathbb{C}}|\overline{S})$ and hence handle $\mathrm{vol}(\mathcal{L}_{i,\mathbb{C}}|\overline{S})$. Need our explicit construction to get fast enough convergence.

A dévissage

Theorem (G'–S.Zhang, 2024)

Assume $g \geq 3$. Then h_{GS} is a dominant height on a Zariski open dense subset $\mathbb{M}_g^{\text{amp}}$ of \mathbb{M}_g defined over \mathbb{Q} . ✓ for (i)



Volume Identity

For each subvariety S of $\mathbb{M}_{g,\mathbb{C}}$,

$$\text{vol}(\widetilde{\mathcal{L}}_{\mathbb{C}}|_S) = \int_{S(\mathbb{C})} c_1(\overline{\mathcal{L}})^{\wedge \dim S}.$$

+

Theorem (GZ 2024)

- $c_1(\overline{\mathcal{L}}) \geq 0$,
- $c_1(\overline{\mathcal{L}})^{\wedge \dim \mathbb{M}_g} \neq 0$ if $g \geq 3$,
- “ $\{x \in S(\mathbb{C}) : (c_1(\overline{\mathcal{L}})|_S^{\wedge \dim S})_x = 0\}$ ” is Zariski closed.



It remains to prove this.

A dévissage

Theorem (G'–S.Zhang, 2024)

Assume $g \geq 3$. Then h_{GS} is a dominant height on a Zariski open dense subset $\mathbb{M}_g^{\text{amp}}$ of \mathbb{M}_g defined over \mathbb{Q} . ✓ for (i)



Volume Identity

For each subvariety S of $\mathbb{M}_{g,\mathbb{C}}$,

$$\text{vol}(\widetilde{\mathcal{L}}_{\mathbb{C}}|_S) = \int_{S(\mathbb{C})} c_1(\overline{\mathcal{L}})^{\wedge \dim S}.$$

+

Theorem (GZ 2024)

- $c_1(\overline{\mathcal{L}}) \geq 0$,
- $c_1(\overline{\mathcal{L}})^{\wedge \dim \mathbb{M}_g} \not\equiv 0$ if $g \geq 3$,
- “ $\{x \in S(\mathbb{C}) : (c_1(\overline{\mathcal{L}})|_S)^{\wedge \dim S}_x = 0\}$ ” is Zariski closed.



It remains to prove this.

Some key ingredients of our proof

Key steps:

- ✓ h_{GS} defined by an a.l.b. $\overline{\mathcal{L}}$
- ✓ volume identity for $\text{vol}(\widetilde{\mathcal{L}})$

Bridged via:

- ✎ Abel–Jacobi periods (Griffiths 1960s)
- ✎ archimedean local heights (Hain 1990s)

- Algebraicity of Betti strata
- Non-vanishing of Betti form

Some tools:

- Adelic line bundle.
- Morse Inequality.
- Mixed Ax–Schanuel (Chiu/Gao–Klingler 2021).
- O-minimality for period map (Bakker, Brunebarbe, Klingler, Tsimerman 2018–2020...).

Some key ingredients of our proof

Key steps:

Main theorem reduced to prove:

- $c_1(\overline{\mathcal{L}}) \geq 0$,
- $c_1(\overline{\mathcal{L}})^{\wedge \dim \mathbb{M}_g} \neq 0$ if $g \geq 3$,
- “ $\{x \in S(\mathbb{C}) : (c_1(\overline{\mathcal{L}})|_S^{\wedge \dim S})_x = 0\}$ ”
is Zariski closed, \forall subvariety S .

Some tools:

- Adelic line bundle.
- Morse Inequality.

Bridged by: (R. de Jong, GZ) $c_1(\overline{\mathcal{L}})$ equals the Betti form.

- Algebraicity of Betti strata
- checkable criterion for non-vanishing of Betti form

- Mixed Ax–Schanuel (Chiu/Gao–Klingler 2021).
- O-minimality for period map (Bakker, Brunebarbe, Klingler, Tsimerman 2018–2020...).

↘ works for any family of homologically trivial cycles.

Some key ingredients of our proof

Key steps:

Main theorem reduced to prove:

- $c_1(\overline{\mathcal{L}}) \geq 0$,
- $c_1(\overline{\mathcal{L}})^{\wedge \dim \mathbb{M}_g} \neq 0$ if $g \geq 3$,
- “ $\{x \in S(\mathbb{C}) : (c_1(\overline{\mathcal{L}})|_S^{\wedge \dim S})_x = 0\}$ ”
is Zariski closed, \forall subvariety S .

Some tools:

- Adelic line bundle.
- Morse Inequality.

Bridged by: (R. de Jong, GZ) $c_1(\overline{\mathcal{L}})$ equals the Betti form.

- Algebraicity of Betti strata
- checkable criterion for non-vanishing of Betti form

- Mixed Ax–Schanuel (Chiu/Gao–Klingler 2021).
- O-minimality for period map (Bakker, Brunebarbe, Klingler, Tsimerman 2018–2020...).

↘ works for any family of homologically trivial cycles.

Some key ingredients of our proof

Key steps:

Main theorem reduced to prove:

- $c_1(\overline{\mathcal{L}}) \geq 0$,
- $c_1(\overline{\mathcal{L}})^{\wedge \dim \mathbb{M}_g} \neq 0$ if $g \geq 3$,
- “ $\{x \in S(\mathbb{C}) : (c_1(\overline{\mathcal{L}})|_S^{\wedge \dim S})_x = 0\}$ ”
is Zariski closed, \forall subvariety S .

Some tools:

- Adelic line bundle.
- Morse Inequality.

Bridged by: (R. de Jong, GZ) $c_1(\overline{\mathcal{L}})$ equals the Betti form.

- Algebraicity of Betti strata
- checkable criterion for non-vanishing of Betti form

- Mixed Ax–Schanuel (Chiu/Gao–Klingler 2021).
- O-minimality for period map (Bakker, Brunebarbe, Klingler, Tsimerman 2018–2020...).

↘ works for any family of homologically trivial cycles.

General setup for studying Betti strata/form/foliation

- $f: X \rightarrow S$ projective morphism over quasi-projective variety, over \mathbb{C} ,
- Z is a family of homologically trivial cycles, of codimension n .

Example (Gross–Schoen and Ceresa)

(GS) $f: \mathcal{C}_g \times_{\mathbb{M}_g} \mathcal{C}_g \times_{\mathbb{M}_g} \mathcal{C}_g \rightarrow \mathbb{M}_g$, Z is the family of Gross–Schoen cycles.
 $n = 2$.

(Ce) $f: \text{Jac}(\mathcal{C}_g/\mathbb{M}_g) \rightarrow \mathbb{M}_g$, Z is the family of Ceresa cycles. $n = g - 1$.

- $\mathbb{V}_Z := Rf_*^{2n-1} \mathbb{Z}_X$. Each fiber $\mathbb{V}_{Z,s} = H^{2n-1}(X_s, \mathbb{Z})$.
- de Rham–Betti comparison $\Rightarrow \mathbb{V}_Z$ is a VHS (variation of Hodge structures) of weight $2n - 1$.
- Polarization (by Lefschetz)

$$Q: \mathbb{V} \otimes \mathbb{V} \rightarrow \mathbb{Q}_S(-n)$$

with $\mathbb{V} := \mathbb{V}_Z \otimes_{\mathbb{Z}_S} \mathbb{Q}_S$.

Intermediate Jacobian and normal function

Definition

The n -th relative intermediate Jacobian is

$$J^n(X/S) := F^n \mathcal{V} \backslash \mathcal{V} / \mathbb{V}_{\mathbb{Z}},$$

with $\mathcal{V} := \mathbb{V}_{\mathbb{Z}} \otimes_{\mathbb{Z}_S} \mathcal{O}_S$ the holomorphic vector bundle.

The fibers are

$$\begin{aligned} (*) \quad J^n(X_S) &= F^n \backslash H^{2n-1}(X_S, \mathbb{C}) / H^{2n-1}(X_S, \mathbb{Z}) && \text{compact complex torus} \\ &\cong H^{2n-1}(X_S, \mathbb{R}) / H^{2n-1}(X_S, \mathbb{Z}) && \text{real torus} \end{aligned}$$

➤ (Griffiths 1969) AJ: $\text{Ch}^n(X_S)_{\text{hom}} \rightarrow J^n(X_S)$.

Definition (Normal function)

$$\nu = \nu_Z: S \rightarrow J^n(X/S), \quad s \mapsto \text{AJ}(Z_s).$$

Betti form, Betti foliation, Betti strata

Family version of (*) becomes

$$J^n(X/S) \xrightarrow{\sim} \mathbb{V}_{\mathbb{R}}/\mathbb{V}_{\mathbb{Z}}.$$

$$\begin{aligned} \nu_{\text{Betti},s}: T_s S &\xrightarrow{d\nu} T_{\nu(s)} J^n(X/S) \\ &\cong T_{\nu(s)} \mathbb{V}_{\mathbb{R}}/\mathbb{V}_{\mathbb{Z}} \\ &= T_s S \oplus \mathbb{V}_{\mathbb{R},s} \rightarrow \mathbb{V}_{\mathbb{R},s} \end{aligned}$$

$\mathbb{V}_{\mathbb{R}}/\mathbb{V}_{\mathbb{Z}} \rightarrow S$ local system of real tori



Betti foliation $\mathcal{F}_{\text{Betti}}$ on $J^n(X/S)$

Definition (Betti form)

$\beta_{\nu}(u, v) := 2Q_s(\nu_{\text{Betti},s}(u), \nu_{\text{Betti},s}(v))$
for all $s \in S(\mathbb{C})$ and $u, v \in T_s S$.

- β_{ν} semi-positive $(1, 1)$ -form (Hain 1990s, using Griffiths' transversality)

Definition (Betti strata)

For any $t \geq 1$, $S^{\text{Betti}}(t) := \{s \in S(\mathbb{C}) : \dim_s(\nu(S) \cap \mathcal{F}_{\text{Betti}}) \geq t\}$.

- $\beta_{\nu}^{\wedge \dim S} \equiv 0 \iff S^{\text{Betti}}(1) = S$

Our result on Betti rank and Betti strata

Theorem (GZ 2024)

- $S^{\text{Betti}}(t)$ is Zariski closed in S .
- We have a checkable criterion for $S^{\text{Betti}}(t) = S$ (equivalently a formula to compute the generic rank of $\nu_{\text{Betti}, S}$). In particular, a checkable criterion for $\beta_v^{\wedge \dim S} \equiv 0$.
- O-minimality for period map to use definable Chow.
- Mixed Ax–Schanuel used [twice](#), second time is through Geometric Zilber–Pink (itself is an application of Ax–Schanuel; [Ullmo](#), Daw–Ren, Gao, Baldi–Klingler–Ullmo, [Baldi–Urbanik](#)).

Back to Gross–Schoen and Ceresa

Main theorem reduced to prove:

For our adelic line bundle $\overline{\mathcal{L}}$ on \mathbb{M}_g with $h_{\overline{\mathcal{L}}} = h_{\text{GS}}$:

- $c_1(\overline{\mathcal{L}}) \geq 0$,
- $c_1(\overline{\mathcal{L}})^{\wedge \dim \mathbb{M}_g} \not\equiv 0$ if $g \geq 3$,
- “ $\{x \in S(\mathbb{C}) : (c_1(\overline{\mathcal{L}})|_S^{\wedge \dim S})_x = 0\}$ ” is Zariski closed, \forall subvariety $S \subseteq \mathbb{M}_{g,\mathbb{C}}$.



(R. de Jong, GZ) $c_1(\overline{\mathcal{L}})$ equals the Betti form β_{GS} .

Corollary (particular case of GZ 2024 on Betti rank and Betti strata)

- $\beta_{\text{GS}} \geq 0$ (Hain 1990s),
- $\beta_{\text{GS}}^{\wedge \dim \mathbb{M}_g} \not\equiv 0$ if $g \geq 3$ (in this case independently by Hain 2024),
- $S(1)$ is Zariski closed, \forall subvariety $S \subseteq \mathbb{M}_{g,\mathbb{C}}$.

Back to Gross–Schoen and Ceresa

Main theorem reduced to prove:

For our adelic line bundle $\overline{\mathcal{L}}$ on \mathbb{M}_g with $h_{\overline{\mathcal{L}}} = h_{\text{GS}}$:

- $c_1(\overline{\mathcal{L}}) \geq 0$,
- $c_1(\overline{\mathcal{L}})^{\wedge \dim \mathbb{M}_g} \not\equiv 0$ if $g \geq 3$,
- “ $\{x \in S(\mathbb{C}) : (c_1(\overline{\mathcal{L}})|_S^{\wedge \dim S})_x = 0\}$ ” is Zariski closed, \forall subvariety $S \subseteq \mathbb{M}_{g,\mathbb{C}}$.



(R. de Jong, GZ) $c_1(\overline{\mathcal{L}})$ equals the Betti form β_{GS} .

Corollary (particular case of GZ 2024 on Betti rank and Betti strata)

- $\beta_{\text{GS}} \geq 0$ (Hain 1990s),
- $\beta_{\text{GS}}^{\wedge \dim \mathbb{M}_g} \not\equiv 0$ if $g \geq 3$ (in this case independently by Hain 2024),
- $S(1)$ is Zariski closed, \forall subvariety $S \subseteq \mathbb{M}_{g,\mathbb{C}}$.

Our main result

Theorem (G'–S.Zhang, 2024)

Assume $g \geq 3$. Let $\mathbb{M}_g^{\text{amp}}$ be the maximal $\overline{\mathbb{Q}}$ -Zariski open subset of \mathbb{M}_g on which h_{GS} is a dominant height.

Then $\mathbb{M}_g^{\text{amp}}$ is non-empty and is defined over \mathbb{Q} . ✓ for (i)

Moreover, $\mathbb{M}_g^{\text{amp}}$ can be “constructed”. ✓ partially for (ii)

Corollary (Generic positivity)

For any number field K , there are at most finitely many C/K lying in $\mathbb{M}_g^{\text{amp}}(\overline{\mathbb{Q}})$ such that $h_{\text{GS}}([C]) \leq 0$.

- **Extra result on torsion:** For every non- $\overline{\mathbb{Q}}$ point $[C]$ in $\mathbb{M}_g^{\text{amp}}$, the cycles $\Delta_{\text{GS}}(C)$ and $\text{Ce}(C)$ are not torsion in the Chow groups. This is also independently proved by Hain (2024) for a non-empty real-analytic open subset and Kerr–Tayou (2024) for a \mathbb{C} -Zariski open dense subset.

Thanks!